

Guillain-Barré Syndrome and Healthcare Needs during Zika Virus Transmission, Puerto Rico, 2016

Emilio Dirlikov, Krista Kniss, Chelsea Major, Dana Thomas, Cesar A. Virgen, Marrielle Mayshack, Jason Asher, Luis Mier-y-Teran-Romero, Jorge L. Salinas, Daniel M. Pastula, Tyler M. Sharp, James Sejvar, Michael A. Johansson, Brenda Rivera-Garcia

To assist with public health preparedness activities, we estimated the number of expected cases of Zika virus in Puerto Rico and associated healthcare needs. Estimated annual incidence is 3.2–5.1 times the baseline, and long-term care needs are predicted to be 3–5 times greater than in years with no Zika virus.

Guillain-Barré syndrome (GBS) is an autoimmune disorder characterized by varying degrees of weakness, sensory abnormalities, and autonomic dysfunction due to peripheral nerve or nerve root damage (1). Annual GBS incidence worldwide is ≈ 1.1 –1.8 cases/100,000 population and varies by geography and age group (2,3). Death is rare and is usually caused by respiratory failure, autonomic dysfunction, or deep vein thrombosis (4).

GBS has been associated with various infectious agents, including Zika virus (5). Zika virus is a flavivirus transmitted primarily by *Aedes* species mosquitoes; symptoms of infection include rash, arthralgia, and fever (6). During a 2013–2014 outbreak in French Polynesia, 42 cases of GBS were reported during a 7-month period, compared with 3–10 cases annually in previous years; all GBS patients during the outbreak had Zika virus antibodies (7).

In December 2015, the Puerto Rico Department of Health reported local transmission of Zika virus (8). In February 2016, the Department of Health reported the first

case of Zika virus–associated GBS and established the GBS Passive Surveillance System, with support from the Centers for Disease Control and Prevention (9). During January 1–July 31, 2016, a total of 56 cases of GBS were reported; evidence of Zika virus or flavivirus infection was found in 34 (61%) of these (9). As in other locations (5), GBS cases in Puerto Rico are anticipated to increase with ongoing Zika virus transmission. To assist with public health preparedness activities, we estimated the annual number of expected cases of GBS and associated healthcare needs in Puerto Rico (online Technical Appendix, <http://wwwnc.cdc.gov/EID/article/23/1/16-1290-Techapp1.pdf>).

The Study

We estimated the weekly number of cases of GBS and associated healthcare needs for 3 scenarios: 1) in the absence of Zika virus transmission; 2) in an average week during Zika virus transmission; and 3) during the peak week of Zika virus transmission (Table). Estimates were derived from baseline and Zika virus–associated GBS cases. The population of Puerto Rico in 2015 was estimated at 3,474,182 persons (10).

We calculated baseline GBS incidence in Puerto Rico by using data collected through medical chart review of patients suspected to have GBS at 9 reference hospitals in Puerto Rico during 2012–2015 and for whom neurologic diagnosis was confirmed by the Brighton Collaboration criteria (11). The 2013 incidence of GBS was 1.7 cases (95% CI 1.3–2.1 cases) per 100,000 population (J.L. Salinas, unpub. data). Using this incidence range, in the absence of Zika virus transmission, we estimated that 1 (interquartile range [IQR] 0–2) case occurs each week, and 59 (IQR 52–66) cases occur each year.

We assumed that, during Zika virus transmission, $\approx 25\%$ of the population could have been infected during 2016, similar to recent chikungunya and dengue virus epidemics in Puerto Rico (12). We used a triangular distribution to characterize uncertainty, with a minimum estimate of 10% infected and a maximum estimate of 70% infected (12,13). Estimated GBS risk associated with Zika virus infection was assumed to be 1.1–2.3 cases/10,000 infections on the basis of a separate analysis of data aggregated from French Polynesia (7), Yap (13), Brazil (14), Colombia, El Salvador, Honduras, the Dominican Republic, and Puerto Rico (L. Mier-y-Teran, unpub. data). We used Monte Carlo sampling to draw 1 million

Author affiliations: Centers for Disease Control and Prevention, Atlanta, Georgia, USA (E. Dirlikov, K. Kniss, C. Major, D. Thomas, M. Mayshack, L. Mier-y-Teran-Romero, J.L. Salinas, D.M. Pastula, T.M. Sharp, J. Sejvar, M.A. Johansson); Puerto Rico Department of Health, San Juan, Puerto Rico (E. Dirlikov, D. Thomas, M. Mayshack, B. Rivera-Garcia); University of California, San Diego, California, USA (C.A. Virgen); Biomedical Advanced Research and Development Authority, Washington, DC, USA (J. Asher); University of Colorado, Denver, Colorado, USA (D.M. Pastula)

DOI: <http://dx.doi.org/10.3201/eid2301.161290>

Table. Estimates of weekly Guillain-Barré syndrome cases and healthcare resource needs, Puerto Rico, 2016

		Estimate		
Variable	Scenario*	Median	Interquartile range	95% Uncertainty interval
New cases and long-term care patients				
Case-patients	No Zika virus	1	0–2	0–4
	Average week during Zika virus	5	3–6	1–11
	Peak week during Zika virus	11	6–17	1–34
Long-term care patients	No Zika virus	0	0–1	0–2
	Average week during Zika virus	2	1–3	0–6
	Peak week during Zika virus	5	2–8	0–16
New patient healthcare resource needs				
Intravenous immunoglobulin	No Zika virus	1	0–2	0–3
	Average week during Zika virus	4	3–6	0–10
	Peak week during Zika virus	10	5–15	0–30
Mechanical ventilation	No Zika virus	0	0–0	0–1
	Average week during Zika virus	1	0–2	0–3
	Peak week during Zika virus	2	1–4	0–8
Regular ward beds	No Zika virus	0	0–1	0–3
	Average week during Zika virus	3	1–4	0–7
	Peak week during Zika virus	6	3–10	0–21
Intensive care unit beds	No Zika virus	0	0–1	0–2
	Average week during Zika virus	2	1–3	0–6
	Peak week during Zika virus	4	2–7	0–15

*The weekly number of Guillain-Barré syndrome cases and associated healthcare needs were estimated for 3 scenarios: 1) in the absence of Zika virus transmission; 2) in an average week during Zika virus transmission; and 3) during the peak week of Zika virus transmission.

values from each of these distributions to estimate each outcome of interest and its associated uncertainty.

We estimated that in an average week of Zika virus transmission, ≈ 5 (IQR 3–6) GBS cases would occur, comprising cases associated with baseline risk and Zika virus infection. As in previous outbreaks of Zika virus and other arboviral diseases, peak weekly incidence could be 2–4 times higher than average incidence. A maximum of 11 (IQR 6–17) cases could occur during the peak week, 2–4 times more than the average number of cases. We predicted that ≈ 241 (IQR 191–305) cases would occur in 2016.

Assumptions regarding treatment needs were also based on data collected from the 2012–2015 Puerto Rico hospitalized patient chart review (J.L. Salinas, unpub. data). We made the following assumptions: 90% of patients will require treatment with intravenous immunoglobulin (IVIg), 20% will require mechanical ventilation, all patients will be hospitalized, 40% will require intensive care, and 45% will require long-term care.

In the absence of Zika virus transmission, the estimated weekly number of new patients treated with IVIg was 1 (IQR 0–2) patient and of those requiring mechanical ventilation was low (IQR 0–0 patients). Estimates for the weekly number of new patients requiring a regular or intensive care unit (ICU) bed were also low (IQR 0–1 patient).

During an average week of Zika virus transmission, ≈ 4 (IQR 3–6) new patients would need treatment with IVIg, and 1 (IQR 0–2) new patient would require mechanical ventilation. An estimated 3 (IQR 1–4) new patients would require a regular ward bed, whereas 2 (IQR 1–3) new patients would require an ICU bed. During the peak week, ≈ 10 (IQR 5–15) new patients would need treatment with

IVIg, and 2 (IQR 1–4) new patients would require mechanical ventilation. An estimated 6 (IQR 3–10) new patients would require a regular ward bed, whereas 4 (IQR 2–7) new patients would require an ICU bed.

An estimated 0 (IQR 0–1) patients would require long-term care during a week without Zika virus transmission, 2 (IQR 1–3) patients during an average week of Zika virus transmission, and 5 (IQR 2–8) patients during the peak week. During 2016, ≈ 108 (IQR 85–138) GBS patients would require long-term care.

Conclusions

We estimated that there would be 191–305 new cases of GBS in Puerto Rico in 2016, comprising baseline and Zika virus-associated cases. This estimate represents an annual incidence of 5.5–8.7 cases/100,000 population, which is 3.2–5.1 times the baseline incidence. Associated healthcare resource needs will increase accordingly. Estimated long-term care needs in 2016 were predicted to be 3–5 times greater than in years with no Zika virus transmission.

These estimates have limitations. First, there is considerable uncertainty around key assumptions, including that increases in GBS incidence will mirror those experienced in other Zika virus-affected countries (5,7,13,14). Second, the estimates of associated healthcare needs did not address all possible needs, such as alternative treatments (i.e., plasmapheresis) or additional treatments, such as those for neuropathic pain, cardiac arrhythmias, and deep vein thrombosis. Third, estimates assumed 1 peak week, although GBS cases tend to cluster, and multiple peaks could occur. Finally, a causal association between Zika virus infection and GBS has not been definitively established.

Continued GBS surveillance will monitor for increased incidence and enable adapted public health response. Healthcare workers, including internists, family physicians, and nurses, might need training to ensure adequate patient clinical management if GBS cases increase as predicted. The Puerto Rico Department of Health and the Centers for Disease Control and Prevention have developed training material toward this end. The availability and accessibility of GBS treatment, especially IVIg, and long-term care services should be evaluated, especially given the high costs of GBS patient care (15). The Puerto Rico Department of Health is also creating an inventory of available and expandable resources, working with manufacturers and distributors to understand supply chains, and facilitating prompt treatment delivery at points of care.

Dr. Dirlikov is a Centers for Disease Control and Prevention Epidemic Intelligence Service Officer posted to the Puerto Rico Department of Health and serves as the GBS team lead for the coordinated Zika response. His areas of interest are infectious diseases, surveillance, and global health.

References

1. Yuki N, Hartung H-P. Guillain-Barré syndrome. *N Engl J Med*. 2012;366:2294–304. <http://dx.doi.org/10.1056/NEJMra1114525>
2. Sejvar JJ, Baughman AL, Wise M, Morgan OW. Population incidence of Guillain-Barré syndrome: a systematic review and meta-analysis. *Neuroepidemiology*. 2011;36:123–33. <http://dx.doi.org/10.1159/000324710>
3. McGrogan A, Madle GC, Seaman HE, de Vries CS. The epidemiology of Guillain-Barré syndrome worldwide. A systematic literature review. *Neuroepidemiology*. 2009;32:150–63. <http://dx.doi.org/10.1159/000184748>
4. Hund EF, Borel CO, Cornblath DR, Hanley DF, McKhann GM. Intensive management and treatment of severe Guillain-Barré syndrome. *Crit Care Med*. 1993;21:433–46. <http://dx.doi.org/10.1097/00003246-199303000-00023>
5. dos Santos T, Rodriguez A, Almiron M, Sanhueza A, Ramon P, de Oliveira WK, et al. Zika virus and the Guillain-Barré syndrome—case series from seven countries. *N Engl J Med*. 2016;NEJMc1609015. <http://dx.doi.org/10.1056/NEJMc1609015>
6. Petersen LR, Jamieson DJ, Powers AM, Honein MA. Zika Virus. *N Engl J Med*. 2016;374:1552–63. <http://dx.doi.org/10.1056/NEJMra1602113>
7. Cao-Lormeau VM, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, et al. Guillain-Barré syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. *Lancet*. 2016;387:1531–9. [http://dx.doi.org/10.1016/S0140-6736\(16\)00562-6](http://dx.doi.org/10.1016/S0140-6736(16)00562-6)
8. Thomas DL, Sharp TM, Torres J, Armstrong PA, Munoz-Jordan J, Ryff KR, et al. Local Transmission of Zika Virus—Puerto Rico, November 23, 2015–January 28, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:154–8. <http://dx.doi.org/10.15585/mmwr.mm6506e2>
9. Dirlikov E, Major CG, Mayshack M, Medina N, Matos D, Ryff KR, et al. Guillain-Barré syndrome during ongoing Zika virus transmission—Puerto Rico, January 1–July 31, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65:910–4. <http://dx.doi.org/10.15585/mmwr.mm6534e1>
10. US Census Bureau. Puerto Rico commonwealth totals: vintage 2015 [cited 2016 May 9]. http://www.census.gov/popest/data/puerto_rico/totals/2015/index.html
11. Sejvar JJ, Kohl KS, Gidudu J, Amato A, Bakshi N, Baxter R, et al.; Brighton Collaboration GBS Working Group. Guillain-Barré syndrome and Fisher syndrome: case definitions and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine*. 2011;29:599–612. <http://dx.doi.org/10.1016/j.vaccine.2010.06.003>
12. Chiu CY, Bres V, Yu G, Krysztof D, Naccache SN, Lee D, et al. Genomic assays for identification of chikungunya virus in blood donors, Puerto Rico, 2014. *Emerg Infect Dis*. 2015;21:1409–13. <http://dx.doi.org/10.3201/eid2108.150458>
13. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med*. 2009;360:2536–43. <http://dx.doi.org/10.1056/NEJMoa0805715>
14. Cardoso CW, Paploski IA, Kikuti M, Rodrigues MS, Silva MM, Campos GS, et al. Outbreak of exanthematous illness associated with Zika, chikungunya, and dengue viruses, Salvador, Brazil. *Emerg Infect Dis*. 2015;21:2274–6. <http://dx.doi.org/10.3201/eid2112.151167>
15. Frenzen PD. Economic cost of Guillain-Barré syndrome in the United States. *Neurology*. 2008;71:21–7. <http://dx.doi.org/10.1212/01.wnl.0000316393.54258.d1>

Address for correspondence: Emilio Dirlikov, Departamento de Salud, Oficina de Epidemiología e Investigación, PO Box 70184, San Juan, PR 00936-8184, USA; email: GBS@salud.pr.gov

Guillain-Barré Syndrome and Healthcare Needs during Zika Virus Transmission, Puerto Rico, 2016

Technical Appendix

Methods

We used Monte Carlo sampling with 1 million realizations to estimate case numbers and resources required. Each realization drew a value of the Guillain-Barré syndrome (GBS) baseline risk, the risk for Zika virus infection, the risk of GBS after Zika virus infection, and the relative incidence at the peak compared with an average week from the distributions specified in the previous section. Once these parameters were selected, the different components of our model were obtained as follows:

Total Number of Baseline and Zika Virus–Associated GBS Cases

The number of baseline GBS cases during the outbreak was sampled from a binomial process for the population of Puerto Rico with a probability equal to the baseline rate, assuming a time span of 1 year. The total number of Zika virus–associated GBS cases during the outbreak was sampled from a binomial process for the population of Puerto Rico with a probability equal to the product of the attack rate and the GBS risk after a Zika virus infection.

Number of Weekly Baseline GBS Cases in the Absence of Zika Virus

The mean number of weekly baseline GBS cases was sampled from a Poisson process with a mean equal to the total number of baseline GBS cases divided by 52, the number of weeks in a year.

Number of Zika Virus–Associated GBS Cases in an Average Week and Peak Week during Zika Virus Transmission

The mean number of weekly Zika virus–associated GBS cases was sampled from a Poisson process with a mean equal to the total number of Zika virus–associated cases divided by

the outbreak duration in weeks (52, the assumed length of the epidemic). The mean number of weekly Zika virus–associated GBS cases in the peak week was sampled by using a Poisson process. The mean was equal to the average number of Zika virus–associated cases per week multiplied by the relative sampled incidence at peak compared with an average week.

Resource Needs for GBS Patients

For each weekly estimate of cases, we estimated the proportion requiring different levels of healthcare from a binomial process with the respective probability (Technical Appendix Table) and the number of estimated cases.

Code

Estimates were generated using R (1). The following code was used:

```
rm(list = ls())

require(data.table)

require(dplyr)

require(triangle)

set.seed(8675309)

n_samples <- 1e6

### fixed parameters

PR_population <- 3474182

GBS_long_term_care_fraction <- 0.45

GBS_IVIG_fraction <- 0.9

GBS_novent_ICU_fraction <- 0.2

GBS_vent_ICU_fraction <- 0.2

GBS_hosp_nonICU_fraction <- 0.6

zika_outbreak_duration_weeks <- 52

### simulation table

sim.data <- data.table(
```

```

GBS_baseline_rate = rnorm(n_samples, 1.7e-5, 0.2e-5),
zika_attack_rate = rtriangle(n_samples, 0.1, 0.7, 0.25),
GBS_zika_rate = rnorm(n_samples, 1.6e-4, 0.31e-4),
peak_mean_weekly_incidence_ratio = runif(n_samples, 2, 4)
) %>%
mutate(
  GBS_total_background_cases =
    rbinom(n_samples, PR_population, GBS_baseline_rate),
  GBS_weekly_background_cases_mean =
    rpois(n_samples, GBS_total_background_cases / zika_outbreak_duration_weeks),
  GBS_total_zika_cases =
    rbinom(n_samples, PR_population, zika_attack_rate * GBS_zika_rate),
  GBS_weekly_zika_cases_mean =
    rpois(n_samples, GBS_total_zika_cases / zika_outbreak_duration_weeks),
  GBS_total_cases = GBS_total_zika_cases + GBS_total_background_cases,
  GBS_weekly_cases_mean = GBS_weekly_zika_cases_mean +
    GBS_weekly_background_cases_mean,
  GBS_weekly_cases_peak = GBS_weekly_background_cases_mean +
    rpois(n_samples, GBS_weekly_zika_cases_mean *
    peak_mean_weekly_incidence_ratio),
  GBS_total_cases_long_term = rbinom(n_samples, GBS_total_cases,
    GBS_long_term_care_fraction),
  GBS_total_cases_background_long_term = rbinom(n_samples,
    GBS_total_background_cases,
    GBS_long_term_care_fraction)

```

```

)

get_resource_needs = function(GBS_weekly_cases) {

  output = rbind(

    data.table(Resource = "Cases per Week,"
    Value = GBS_weekly_cases),

    data.table(Resource = "Long-Term Care Patients,"
    Value = rbinom(n_samples, GBS_weekly_cases, GBS_long_term_care_fraction)),

    data.table(Resource = "Ventilators,"
    Value = rbinom(n_samples, GBS_weekly_cases, GBS_vent_ICU_fraction)),

    data.table(Resource = "IVIg per Week,"
    Value = rbinom(n_samples, GBS_weekly_cases, GBS_IVIG_fraction)),

    data.table(Resource = "Non-ICU Beds,"
    Value = rbinom(n_samples, GBS_weekly_cases, GBS_hosp_nonICU_fraction)),

    data.table(Resource = "ICU Beds,"
    Value = rbinom(n_samples, GBS_weekly_cases,
    GBS_novent_ICU_fraction + GBS_vent_ICU_fraction)))

  return(output)

}

raw_weekly_resources <- rbind(

  cbind(get_resource_needs(sim.data$GBS_weekly_background_cases_mean),
  Estimate = "No Zika"),

  cbind(get_resource_needs(sim.data$GBS_weekly_cases_mean),
  Estimate = "Average Zika"),

  cbind(get_resource_needs(sim.data$GBS_weekly_cases_peak),

```

```

Estimate = "Peak Zika")) %>%

mutate(

Estimate = factor(Estimate, c("No Zika," "Average Zika," "Peak Zika")),

Resource = factor(Resource, c("Cases per Week," "Long-Term Care Patients,"

"IVIg per Week," "Ventilators," "Non-ICU Beds," "ICU Beds"))))

### summarize

summarized_data <- raw_weekly_resources %>%

as.data.frame() %>%

group_by(Resource, Estimate) %>%

do({

quantile_frame =

data.frame(t(quantile(.$Value, probs = c(0.025, 0.25, 0.5, 0.75, 0.975))))

colnames(quantile_frame) = c("Low," "IQR_Low," "Mid," "IQR_High," "High")

return(quantile_frame)

}) %>%

mutate(

Median = signif(Mid, 2),

IQR = paste0(signif(IQR_Low, 2), "-", signif(IQR_High, 2)),

`95% UT` = paste0(signif(Low, 2), "-", signif(High, 2))) %>%

select(Resource, Estimate, Median, IQR, `95% UT`) %>%

arrange(Resource, Estimate)

summarized_data[]

```


References

1. R Project. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2016.
2. US Census Bureau. Puerto Rico commonwealth totals: vintage 2015 [cited 2016 May 9].
http://www.census.gov/popest/data/puerto_rico/totals/2015/index.html
3. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med*. 2009;360:2536–43.
<http://dx.doi.org/10.1056/NEJMoa0805715>
4. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med*. 2009;360:2536–43.
<http://dx.doi.org/10.1056/NEJMoa0805715>
5. Cardoso CW, Paploski IA, Kikuti M, Rodrigues MS, Silva MM, Campos GS, et al. Outbreak of exanthematous illness associated with Zika, chikungunya, and dengue viruses, Salvador, Brazil. *Emerg Infect Dis*. 2015;21:2274–6. <http://dx.doi.org/10.3201/eid2112.151167>

Technical Appendix Table. Parameters used in the model for a study of Guillain-Barré syndrome and healthcare needs during Zika virus transmission, Puerto Rico, 2016*

Parameter	Estimated value	Distribution	Source	Citation
Puerto Rico population, 2016	3,474,182	Fixed	US Census	US Census (2)
Baseline GBS rate	1.7/100,000 population/year	Gaussian, $\sigma = 0.20$	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Zika virus infection risk in Puerto Rico, 2016	25% peak probability	Triangle, min 10%, max 70%	Historic CHIKV and DENV epidemics in Puerto Rico	(3)
Fraction of Zika virus infections developing GBS	1.6/10,000 infections	Gaussian, $\sigma = 0.31$	Analysis of country-level GBS and Zika virus reports	(L. Mier-y-Teran, unpub. data) (3–5)
Relative Zika virus incidence during peak week compared with an average week	2–4 times	Uniform	Analysis of country-level GBS and Zika virus reports	(L. Mier-y-Teran, unpub. data)
Percentage of GBS patients needing long-term care	45	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Percentage of GBS patients needing IVIg	90	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Percentage of GBS patients needing an ICU bed	40	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Percentage of GBS patients needing an ICU bed and a ventilator	20	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Percentage of GBS patients needing an ICU bed but no ventilator	20	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)
Percentage of GBS patients needing a regular ward bed	60	Fixed	Chart review of Puerto Rico GBS patients	(J.L. Salinas, unpub. data)

*CHIKV, chikungunya virus; DENV, dengue virus; GBS, Guillain-Barré syndrome; ICU, intensive care unit; IVIg, intravenous immunoglobulin.